

Randomized trial of dietary fiber and *Lactobacillus casei* administration for prevention of colorectal tumors

Ishikawa H¹, Akedo I², Otani T³, Suzuki T⁴, Nakamura T¹, Takeyama I¹, Ishiguro S⁵, Miyaoka E⁶, Sobue T⁷, Kakizoe T⁸

¹ *Laboratory of Hereditary Tumors, Institute for Advanced Medical Sciences, Hyogo College of Medicine, Osaka*

² *Internal Medicine, Osaka Central Hospital, Osaka*

³ *Department of Gastroenterology, Osaka Medical Center for Cancer and Cardiovascular Diseases, Osaka*

⁴ *Department of Cancer Epidemiology, Osaka Medical Center for Cancer and Cardiovascular Diseases, Osaka*

⁵ *Department of Pathology, Osaka Medical Center for Cancer and Cardiovascular Diseases, Osaka*

⁶ *Department of Mathematics, Tokyo University of Science, Tokyo*

⁷ *Cancer Information and Epidemiology Division, Research Institute, National Cancer Center, Tokyo*

⁸ *President, National Cancer Center, Tokyo*

The epidemiologic evidence that dietary fiber protects against colorectal cancer is equivocal. No large-scale clinical study of the administration of *Lactobacillus casei* has been reported. We examined whether dietary fiber and *L. casei* prevented the occurrence of colorectal tumors. Subjects were 398 men and women presently free from tumor who had had at least 2 colorectal tumors removed. Subjects were randomly assigned to 4 groups administered wheat bran, *L. casei*, both or neither. The primary end point was the presence or absence of new colorectal tumor(s) diagnosed by colonoscopy after 2 and 4 years. Among 380 subjects who completed the study, 95, 96, 96 and 93 were assigned to the wheat bran, *L. casei*, both and no treatment groups, respectively. Multivariate adjusted ORs for occurrence of tumors were 1.31 (95% CI 0.87–1.98) in the wheat bran group and 0.76 (0.50–1.15) in the *L. casei* group compared to the control group. There was a significantly higher number of large tumors after 4 years in the wheat bran group. The occurrence rate of tumors with a grade of moderate atypia or higher was significantly lower in the group administered *L. casei*. No significant difference in the development of new colorectal tumors was observed with administration of either wheat bran or *L. casei*. However, our results suggest that *L. casei* prevented atypia of colorectal tumors.